

Constraining dark matter through 21 cm observations

M. Valdés¹, A. Ferrara¹, M. Mapelli¹, E. Ripamonti²

¹ *SISSA/ISAS, via Beirut 2-4, 34014 Trieste, Italy;*

² *Kapteyn Astronomical Institute, University of Groningen, Postbus 800, 9700 AV, Groningen, The Netherlands.*

5 February 2008

ABSTRACT

Beyond reionization epoch cosmic hydrogen is neutral and can be directly observed through its 21 cm line signal. If dark matter (DM) decays or annihilates the corresponding energy input affects the hydrogen kinetic temperature and ionized fraction, and contributes to the Ly α background. The changes induced by these processes on the 21 cm signal can then be used to constrain the proposed DM candidates, among which we select the three most popular ones: (i) 25-keV decaying sterile neutrinos, (ii) 10-MeV decaying light dark matter (LDM) and (iii) 10-MeV annihilating LDM. Although we find that the DM effects are considerably smaller than found by previous studies (due to a more physical description of the energy transfer from DM to the gas), we conclude that combined observations of the 21 cm background and of its gradient should be able to put constraints at least on LDM candidates. In fact, LDM decays (annihilations) induce differential brightness temperature variations with respect to the non decaying/annihilating DM case up to $\Delta\delta T_b = 8$ (22) mK at about 50 (15) MHz. In principle this signal could be detected both by current single dish radio telescopes and future facilities as LOFAR; however, this assumes that ionospheric, interference and foreground issues can be properly taken care of.

Key words: intergalactic medium - cosmology: theory - diffuse radiation - dark matter

1 INTRODUCTION

Cosmic recombination at $z \sim 1000$ left the gas in the Universe in a nearly uniform, dark, neutral state which we currently denote as the Dark Ages of the Universe. Gas remained neutral until the first luminous objects emerged at $z \sim 20$ –30, leading to the complete reionization of the intergalactic medium (IGM) at a later, yet unknown, epoch.

The investigation of the Dark Ages is one of the frontiers of modern cosmology and a new generation of radio interferometers is currently in development to study the redshifted 21 cm hyperfine triplet-singlet level transition of the ground state of neutral hydrogen. Instruments such as the LOw Frequency ARray (LOFAR), the 21 Centimeter Array (21CMA), the Mileura Wide-field Array (MWA) and the Square Kilometer Array (SKA), are expected to reach the sensitivity required to map the HI distribution at angular resolution of the order of a few arcminutes (e.g. Pen, Wu, & Peterson 2004; Bowman, Morales, & Hewitt 2005; Kassim et al. 2004; Wyithe, Loeb, & Barnes 2005).

The main scientific goal of such instruments is to perform an accurate tomography of matter during these remote cosmic epochs, and reconstruct in detail the latest phases of the reionization history. However, this might not be the only major achievement of such experiments. For example,

an analysis of the features of HI 21 cm maps from the Dark Ages could also allow to put constraints on the existence and nature of decaying and annihilating Dark Matter (DM). In fact, if DM actually decays or annihilates, the injection of high energy photons into the IGM would heat and ionize the neutral gas, leaving an imprint on the 21 cm background signal which could be directly observed (Shchekinov & Vasiliev 2006; Furlanetto, Oh, & Pierpaoli 2006).

The importance of DM decays and annihilations on the reionization history has been analyzed in detail by several authors. Most authors have come to the conclusion that the effects of DM on the overall reionization process are relatively small and they cannot compete with those induced by more conventional ionizing sources as stars and (mini-) quasars. However, as we will explain in detail later on, radiation from decaying/annihilating DM might change the thermal and ionization history of the gas to such an extent that differences with the case in which DM is not considered are large enough to produce a clear signature in observable 21 cm signal. Assessing the amplitude of these deviations from the standard scenario requires a careful study of complex physical processes, and in particular of the amount of energy that can be transferred from the decay/annihilation products to the IGM. A recent study (Ripamonti, Mapelli, & Ferrara 2006a, hereafter RMF06a) finds that only a rela-

tively small fraction of the injected energy is effectively absorbed by the IGM and goes into heating and reionization. As a result, the effects on the 21 cm background signal might be smaller than previously predicted (Shchekinov & Vasiliev 2006; Furlanetto et al. 2006); hence, the question remains if the DM signal can still be observed by next generation radio interferometers. If so, this experiment would constitute a superb tool to distinguish among different DM candidates via their decaying/annihilating properties. In this paper we calculate the effects of DM decays/annihilations on the 21 cm background for some of the most popular DM candidates, such as decaying or annihilating Light Dark Matter (LDM) and decaying sterile neutrinos.

The rest of the paper is organized as follows. In Sec. 2 we briefly introduce the DM candidates and their effect on the IGM; in Sec. 3 we provide the basic equations to study the 21 cm radiation in presence of decaying/annihilating DM. In Sec. 4 we present the results of our calculations, which are then discussed in Sec. 5. Throughout the paper we assume, in agreement with the 3-yr WMAP data analysis (Spergel et al. 2006), a Λ CDM cosmology with $\Omega_m = 0.24$, $\Omega_\Lambda = 0.76$, $h = 0.73$, $\Omega_b = 0.042$, $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 BASIC PHYSICS

We are interested in calculating the effects on HI 21 cm line signal produced by two among the most popular low-mass DM candidates, *i.e.* LDM ($\sim 1 - 10 \text{ MeV}$) and sterile neutrinos ($\sim 2 - 50 \text{ keV}$). In the case of sterile neutrinos only the decay process is allowed; LDM particles instead can both decay and annihilate. We neglect heavier DM candidates (with mass larger than $\sim 100 \text{ MeV}$) because previous studies (Mapelli, Ferrara & Pierpaoli 2006) have already shown that, even assuming that all the energy released following annihilations is immediately absorbed by the IGM, they represent a negligible heating/ionization source for the gas.

Both in the case of sterile neutrinos and of LDM, the rate of energy transfer per baryon to the IGM can be written as (RMF06a; Ripamonti, Mapelli & Ferrara 2006b, hereafter RMF06b):

$$\dot{E}_x(z) = f_{abs}(z) \dot{n}_{DM}(z) m_{DM} c^2, \quad (1)$$

where m_{DM} is the mass of the DM particle and c is the speed of light. The energy absorbed fraction, $f_{abs}(z)$, is discussed in detail in the following; $\dot{n}_{DM}(z)$ is the decrease rate of the number of DM particles per baryon.

In the case of DM decays, $\dot{n}_{DM}(z)$ is given by

$$\dot{n}_{DM}(z) \simeq \frac{n_{DM,0}}{\tau_{DM}}, \quad (2)$$

where $n_{DM,0}$ and τ_{DM} are the current number of DM particles per baryon and the lifetime of DM particles, respectively.

For the annihilations:

$$\dot{n}_{DM}(z) \simeq \frac{1}{2} n_{DM,0}^2 \mathcal{N}_b(0) \langle \sigma v \rangle (1+z)^3, \quad (3)$$

where $\mathcal{N}_b(0) = 2.5 \times 10^{-7} \text{ cm}^{-3}$ is the current baryon number density (Spergel et al. 2006), and $\langle \sigma v \rangle$ is the thermally-averaged DM annihilation cross section. The values of m_{DM} ,

$n_{DM,0}$, τ_{DM} and $\langle \sigma v \rangle$ depend from the properties of the chosen DM candidate (see Secs. 2.1 and 2.3).

2.1 The energy absorbed fraction

The most delicate term in eq. (1) is represented by $f_{abs}(z)$, *i.e.* the fraction of the DM particle rest mass that is absorbed by the gas at a given redshift z ; f_{abs} strongly depends on how the decay/annihilation products interact with the IGM. Since accounting for the physical processes which govern such interactions is quite complicated, most of previous studies assume that: (i) all the energy released by DM decays/annihilations is immediately absorbed (Hansen & Haiman 2004; Pierpaoli 2004; Biermann & Kusenko 2006; Mapelli et al. 2006), (ii) leave f_{abs} as a free parameter (Padmanabhan & Finkbeiner 2005; Zhang et al. 2006), or (iii) make a partial treatment of the energy redistribution (Chen & Kamionkowski 2004; Mapelli & Ferrara 2005).

Recently, RMF06a have calculated behaviour of $f_{abs}(z)$ in detail, for the most common case in which the decay/annihilation products are photons, active neutrinos or electron-positron pairs. Photons are affected by Compton scattering and photo-ionization; for pairs, the relevant processes are inverse Compton scattering, collisional ionizations, and positron annihilations.

RMF06a found that, if the decay/annihilation products are either photons and active neutrinos or pairs, f_{abs} is close to the maximum allowed value (equal to 0.5 for sterile neutrinos, due to the active neutrino production, and equal to 1 for LDM) only at very high redshift ($z \gg 100$). At lower redshifts, f_{abs} rapidly drops to values < 0.1 ; also, the higher is the mass of the progenitor DM particle, the faster is the decrease of f_{abs} . Thus, accounting for the correct $f_{abs}(z)$ determination dramatically reduces the possible effects of DM decays and annihilations on the IGM heating and ionization (RMF06a) with respect to previous estimates.

In this paper, for the first time, we will adopt the correct estimate of $f_{abs}(z)$ (as given in RMF06a) in order to evaluate the impact of DM decays and annihilations on 21 cm emission. Previous papers (e.g. Furlanetto et al. 2006) have assumed a redshift-independent $f_{abs}(z)$, which appears to be quite unrealistic (for a discussion, see Sec. 4) and leads to optimistic upper limits for the contribution of DM-related effects to 21 cm maps.

Differently from Furlanetto et al. 2006, who do not select any specific DM candidate and leave the decay rate as a free parameter, we chose to consider three specific DM candidates (sterile neutrinos, decaying LDM and annihilating LDM). This choice allows us to give predictions which can be more easily related to other DM measurements, such as the X-ray constraints on the sterile neutrino mass (Watson et al. 2006; Boyarsky et al. 2006a) or the detection of the 511-keV emission line from the Galactic center (Knödseder et al. 2005).

2.2 Sterile neutrinos

Many models of sterile neutrinos have been proposed, with mass ranging from eV to TeV. Here, we are interested in sterile neutrinos as warm DM (WDM) candidates, with masses of the order of a few keV ($\sim 2 - 50 \text{ keV}$). These particles

can decay into an active neutrino and a photon (Dolgov 2002 and references therein).

The mass (and thus the lifetime) of radiatively decaying sterile neutrinos can be constrained by the absence of any detection of X-ray lines consistent with photons due to sterile neutrino decays in galaxy clusters (Abazajian, Fuller & Tucker 2001; Abazajian 2006; Abazajian & Koushiappas 2006; Boyarsky et al. 2006b) or in galaxies (Watson et al. 2006 obtained the strongest constraints from the study of the Andromeda galaxy). Other constraints come from the comparison between the unresolved X-ray background and the expected contribution from sterile neutrino decays (Mapelli & Ferrara 2005; Boyarsky et al. 2006a).

In this paper we consider the representative case of $m_\nu = 25$ keV sterile neutrinos, whose contribution to heating is maximum (RMF06a), due to the weakness of the available constraints on lifetime for neutrinos of such mass. For such particle the upper limits on the lifetime and the present number density number per baryon are $\tau_{\text{DM}} = 9.67 \times 10^{25}$ s and $n_{\text{DM},0} = 1.88 \times 10^5$, respectively (see RMF06a, RMF06b). The contribution of other sterile neutrinos masses to the 21 cm line is expected to be comparable or smaller than for the case considered here.

2.3 Light dark matter

We define as LDM particles all the DM candidates whose mass is $1 \leq m_{\text{LDM}}/\text{MeV} \leq 100$. Such particles have been suggested as a possible source for the detected 511-keV excess from the Galactic centre (Knödseder et al. 2005). According to this scenario, their maximum allowed mass m_{LDM} should be 20 MeV, not to overproduce detectable gamma rays via internal bremsstrahlung (Beacom, Bell & Bertone 2004), or even ~ 3 MeV, if we consider also the production of gamma rays for inflight annihilations of the positrons (Beacom & Yüksel 2006).

LDM can decay or annihilate, producing photons, neutrinos and pairs. We will treat both channels in detail and assume that the only decay/annihilation products are pairs, an assumption leading to an upper limit in terms of IGM heating (RMF06a).

We consider, as a template, the case of 10-MeV LDM particles, which again yields the most efficient heating case (see RMF06a). For such particles the upper limits of the current number (per baryon), the lifetime and the cross-section are $n_{\text{DM},0} \sim 446$, $\tau_{\text{DM}} = 4 \times 10^{25}$ s, and $\langle \sigma v \rangle \sim 2.4 \times 10^{-28} \text{ cm}^3 \text{ s}^{-1}$, respectively (RMF06a).

3 EFFECTS ON THE 21 CM RADIATION

3.1 Ly α pumping

The 21 cm line is associated with the hyperfine transition between the triplet and the singlet levels of the hydrogen ground state. This transition is governed by the spin temperature, T_S , defined as:

$$\frac{n_1}{n_0} = 3 \exp\left(-\frac{T_\star}{T_S}\right), \quad (4)$$

where n_0 and n_1 are the number densities of hydrogen atoms in the singlet and triplet ground hyperfine levels, and $T_\star =$

0.068 K is the temperature corresponding to the transition energy.

In the presence of the Cosmic Microwave Background (CMB) alone, the spin temperature reaches thermal equilibrium with $T_{\text{CMB}} = 2.73(1+z)$ K on a short time-scale, making the HI undetectable in emission or absorption.

Two mechanisms can decouple T_S from T_{CMB} : (i) collisions, which are effective mainly at $z \geq 70$ due to the higher mean IGM density, and (ii) scattering by Ly α photons – the so-called Wouthuysen-Field process or Ly α pumping (e.g. Wouthuysen 1952; Field 1959; Hirata 2005) – which couples T_S to the kinetic gas temperature T_K via the mixing of the hyperfine levels of the HI ground state through intermediate transitions to the excited $2p$ state.

The spin temperature is then given by the equation:

$$T_S = \frac{T_{\text{CMB}} + y_\alpha T_k + y_c T_k}{1 + y_\alpha + y_c} \quad (5)$$

where T_k is the kinetic temperature, and the Ly α and collisional coupling coefficients are given by the following expressions:

$$y_\alpha = \frac{P_{10} T_\star}{A_{10} T_k}, \quad (6)$$

$$y_c = \frac{C_{10} T_\star}{A_{10} T_k}, \quad (7)$$

where

$$C_{10} = k_{10} n_{\text{HI}} + n_e \gamma_e, \quad (8)$$

and

$$P_{10} = \frac{16}{27} \frac{\pi J_\alpha \sigma_\alpha}{h_p \nu_\alpha}. \quad (9)$$

In the above equations $A_{10} = 2.85 \times 10^{-15} \text{ s}^{-1}$ is the spontaneous emission coefficient of the 21 cm line, P_{10} is the indirect de-excitation rate of the hyperfine structure levels. We write $P_{10} = 4P_\alpha/27$; in addition, $P_\alpha = (4\pi J_\alpha \sigma_\alpha)/(h_p \nu_\alpha)$ is the outcome of the equation for the Ly α scattering rate,

$$P_\alpha = c \int d\nu n(\nu) \frac{\sigma(\nu)}{\nu} = \frac{4\pi}{h} \int d\nu \frac{J_\nu \sigma(\nu)}{\nu}, \quad (10)$$

in the case that $\sigma(\nu)$ is a δ function. A detailed investigation of the physics of the Wouthuysen-Field process (Hirata 2005) finds small corrections to these expressions which we neglect here for simplicity. The coefficient C_{10} in eq. (8) is the collisional de-excitation rate by hydrogen atoms and electrons, where k_{10} is tabulated in Allison & Dalgarno (1969) for different temperatures; γ_e , according to Liszt (2001), is

$$\log \gamma_e(T_k) = -9.607 + \log T_k^{1/2} \exp(-(\log T_k)^{4.5}/1800). \quad (11)$$

for $T_k \leq 10^4$ K, while $\gamma_e(T_k > 10^4 \text{ K}) = \gamma_e(T_k = 10^4 \text{ K})$.

For the Ly α pumping to be effective, a minimum Ly α background intensity, J_α , is required. This is given by the condition (Ciardi & Madau 2003):

$$J_\alpha \geq 9 \times 10^{-23} (1+z) \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}, \quad (12)$$

at the redshift of interest.

3.2 IGM evolution and 21 cm background.

Next, we want to understand how DM decays/annihilations affect the thermal and ionization evolution of the IGM,




Figure 1. *Left panel:* T_S and T_k as a function of redshift (25-keV sterile neutrino decays, f_{abs} as in RMF06a). Thin solid line: T_{CMB} ; thick solid (short-dashed) line: T_S with (without) 25-keV neutrino decays; dotted (long-dashed) line: T_k with (without) 25-keV neutrino decays. *Right:* T_S and T_k as a function of redshift for 25-keV sterile neutrino decays and for $f_{abs} = 0.5$. The lines are the same as in the left panel.

which, in turn, determines the level of the 21 cm background signal. The equation that describes the evolution of the ionized fraction x_e is the following (see e.g. Chen & Kamionkowski 2004):

$$-\frac{dx_e}{dz} = \frac{1}{H(z)(1+z)}[R_s(z) - I_s(z) - I_x(z)], \quad (13)$$

where $I_x = \dot{E}_x/E_0$ is the contribution to the ionization rate due to DM; I_s and R_s are the standard ionization and recombination rates per baryon. Considering eqs. (1)-(3) we have that

$$I_x = \chi_i(z) \frac{\dot{E}_x}{E_0} = f_{abs}(z) \chi_i(z) \Gamma_x f_x \frac{m_p c^2}{E_0} \quad (14)$$

for DM decays, and

$$I_x = \chi_i(z) \frac{\dot{E}_x}{E_0} = f_{abs}(z) \chi_i(z) f_x \frac{m_p c^2}{E_0} n_{DM,0} \mathcal{N}_b(z) \langle \sigma v \rangle \quad (15)$$

for DM annihilations. In the last two equations $E_0 = 13.6$ eV is the hydrogen ionization threshold, m_p is the proton mass, χ_i is the fraction of the energy absorbed by the IGM from DM decays/annihilations that goes into ionizations ($\chi_i \sim (1 - x_e)/3$, see Shull 1979, Shull & van Steenberg 1985), $f_x = \Omega_x(z)/\Omega_b(z)$, $\mathcal{N}_b(z)$ is the baryon number density and $\Gamma_x = 1/\tau_{DM}$.

The equation regulating the evolution of IGM temperature can be written as:

$$(1+z) \frac{dT_k}{dz} = 2T_k + \frac{l_\gamma x_e}{H(z)(1+f_{He}+x_e)} (T_k - T_{CMB}) - \frac{2\chi_h \dot{E}_x}{3k_b H(z)(1+f_{He}+x_e)} \quad (16)$$

where $l_\gamma = (8\sigma_T a_R T_{CMB}^4)/(3m_e c)$, χ_h is the fraction of the

absorbed DM energy deposited into the IGM as heating (Shull & van Steenberg 1985) and $f_{He} = 0.24$ is the helium fraction by mass. The Ly α heating resulting from repeated scatterings as the photons are redshifted into the Lyman resonances is negligible as it has been recently shown (see e.g. Chen & Miralda Escudé 2004). The ionization and temperature eqs. (13)-(16) are solved using a modified version of RECFAST (Seager et al. 1999).

A third equation is needed in order to compute the 21 cm background: the one describing the evolution of the Ly α background intensity J_α . Following Madau, Meiksin & Rees (1997) we write:

$$J_\alpha(z) = \frac{\mathcal{N}_H^2 hc}{4\pi H(z)} \left[x_e x_p \alpha_{2^2P}^{eff} + x_e x_{HI} \gamma_{eH} + \frac{\chi_\alpha \dot{E}_x(z)}{\mathcal{N}_H h \nu_\alpha} \right], \quad (17)$$

where the first two terms are the contributions from recombinations and collisional excitations by electron impacts, while the third term is the DM contribution.

In the last equation $\mathcal{N}_H = 0.92 \mathcal{N}_b$ is the number density of hydrogen atoms, $\alpha_{2^2P}^{eff}$ is the effective recombination coefficient to the 2^2P level (Pengelly 1964), which includes direct recombinations to the 2^2P level and recombinations to higher levels, followed by transitions to the 2^2P level via all possible cascade paths.

Finally, $\gamma_{eH} \approx 2.2 \times 10^{-8} \exp(-11.84/T_4) \text{ cm}^3 \text{ s}^{-1}$ is the collisional excitation rate of HI atoms photons by electron impacts; χ_α is the net fraction of the absorbed X-ray photons from DM decays/annihilations which is converted into Ly α photons (Shull & van Steenberg 1985), and $T_4 = T/(10^4 \text{ K})$.

Once $T_S(z)$ has been determined through eq. (5), we can obtain the 21 cm radiation intensity, which can be expressed




Figure 2. *Left panel:* T_S and T_k as a function of redshift (10-MeV LDM decays, f_{abs} as in RMF06a). Thin solid line: T_{CMB} ; thick solid (short-dashed) line: T_S with (without) 10-MeV LDM decays; dotted (long-dashed) line: T_k with (without) 10-MeV LDM decays. *Right:* T_S and T_k as a function of redshift for 10-MeV LDM decays and for $f_{abs} = 1$. The lines are the same as in the left panel.

by the differential brightness temperature between a neutral hydrogen patch and the CMB:

$$\delta T_b \simeq \frac{T_S - T_{CMB}}{1 + z} \tau, \quad (18)$$

where τ is the optical depth of the neutral IGM at $21(1+z)$ cm:

$$\tau \simeq \frac{3c^3 h_p A_{10}}{32\pi k_B \nu_0^2 T_S H(z)} \mathcal{N}_{\text{HI}}. \quad (19)$$

In equation (19), h_p and k_B are the Planck and Boltzmann constants, respectively, $\nu_0 = 1420$ MHz is the 21 cm hyperfine transition frequency, and $\mathcal{N}_{\text{HI}}$ is the local HI number density. If T_S is higher than T_{CMB} , the neutral IGM will be visible in emission against the CMB; on the contrary, if $T_S < T_{CMB}$ it will be visible in absorption.

4 RESULTS

In a Universe where DM does not decay or annihilate the spin temperature and the kinetic temperature of the gas track the CMB temperature down to $z \approx 300$, when the kinetic temperature starts to decrease adiabatically, $T_K \propto (1+z)^2$, while $T_{CMB} \propto (1+z)$. The collisions at this redshift are efficient at coupling T_S and T_K : the spin temperature subsequently tracks the kinetic temperature down to $z \sim 70$. At lower redshifts radiative coupling to the CMB becomes dominant and $T_S \rightarrow T_{CMB}$ again. As a result, for $30 \leq z \leq 300$ HI is visible in absorption against the CMB at wavelength of $21(1+z)$ cm. This scenario could change considerably if we allow for DM decay/annihilation, depressing or even erasing the absorption feature discussed above. Also, we note that different DM candidates leave different traces on the 21 cm background signal, and therefore it could be

possible, in principle, to constrain directly the DM nature through 21 cm observations. To isolate the effect of DM, we will assume in the following that the entire cosmic dark matter content is constituted by particles of the considered type (sterile neutrinos or LDM), *i.e.* $\Omega_X = \Omega_m - \Omega_b$.

4.1 WDM: 25-keV sterile neutrino decay

In Fig. 1 we compare the effects of 25-keV sterile neutrino decays on T_S assuming either the physically motivated value of f_{abs} (left panel) or the commonly adopted value $f_{abs} = 0.5$ (right panel), *i.e.* the maximum allowed value for sterile neutrinos corresponding to complete absorption. In addition, in each of the panels we explore the differences between models with or without the energy injection term due to DM decays. We will refer in the following to the latter as the *standard case*, for brevity.

With respect to the case without DM energy injection, we find, independently of the assumption made for f_{abs} , a higher kinetic temperature and hydrogen ionization fraction. However, such an enhancement produces only very modest T_S differences with respect to the standard case, which has some effect on the differential brightness temperature δT_b . The difference $T_S - T_{CMB}$ remains extremely small due to the limited ability of the Ly α pumping to decouple the two temperatures.

The main difference between the results obtained from the two f_{abs} assumptions is the behavior of T_K at $z \sim 10-70$. When we use the realistic f_{abs} value, the IGM kinetic temperature positively deviates at most by 50% from its standard evolution; if instead $f_{abs} = 0.5$ is adopted (e.g. as assumed for example by Furlanetto et al. 2006), the kinetic temperature overshoots the CMB one and reaches very large values (> 100 K) at low redshifts.




Figure 3. *Left panel:* T_S and T_K as a function of redshift (10-MeV LDM annihilations, f_{abs} as in RMF06a). Thin solid line: T_{CMB} ; thick solid (short-dashed) line: T_S with (without) 10-MeV LDM annihilations; dotted (long-dashed) line: T_K with (without) LDM annihilations. *Right:* T_S and T_K as a function of redshift for 10-MeV LDM annihilations and for $f_{abs} = 1$. The lines are the same as in the left panel.

4.2 LDM: 10-MeV decay/annihilation

We now turn to the analysis of LDM candidates. Following the same procedure as in the previous Section we compare the results of the two different assumptions for f_{abs} (note that for LDM, the constant absorbed fraction case corresponds to $f_{abs} = 1$, see Sec. 2.1), and also test the effects of decays/annihilation against the standard case.

We start from the analysis of the realistic f_{abs} case (Fig. 2, left panel) for decaying LDM. The most striking feature is that the evolution of T_K decouples from the adiabatic one already at $z \approx 50$, and starts to increase below $z \approx 20$, reaching $T_K = 30$ K at $z = 10$. Such thermal history forces T_S to remain below T_{CMB} for a much longer redshift interval (down to $z = 10$) than in the standard case, in which $T_S \approx T_{CMB}$ already at $z \approx 25$. Interestingly, this effect extends the frequency range in which the IGM can be observed in absorption to higher values. The main physical reason for the larger impact of LDM decays on T_S with respect to sterile neutrinos analyzed above basically lies in the larger f_{abs} value (approximately a factor of 10 below $z = 30$) of LDM particles; this is on top of their larger rest mass. The higher heating rate increases both T_K and the Ly α background (*i.e.* y_α), thus pushing T_S away from T_{CMB} .

When f_{abs} is instead artificially forced to be constant and equal to unity (Fig. 2, right panel), the physical arguments given above can still be used to interpret the results, but the deviation from the standard case is much more dramatic, and results in an overestimate of the LDM decays impact. For example, T_K becomes larger than T_{CMB} already at $z \approx 50$ and stays above it thereafter, reaching > 1000 K at $z = 10$. The increase is so strong that it essentially erases the absorption feature expected in the standard case above $z = 30$. The IGM according to this prescription should be observed in emission up to very high redshifts.

For comparison sake, we note that this case has nearly the same decay rate, $\Gamma_X = 2.5 \times 10^{-26} \text{ s}^{-1}$, as that assumed in one of the cases explored by Furlanetto et al. (2006) (dot-dashed line in Fig. 1 of that paper). Consistently, the evolution of T_K from the two studies is essentially the same.

Finally we turn to the case of annihilating LDM (Fig. 3). As usual, the left panel of that figure reports the case in which the realistic, redshift dependent, values of f_{abs} are assumed. In this case, the kinetic temperature deviates from the standard adiabatic evolution at extremely high redshifts ($z > 200$): this is because the annihilation process depends on the square of the baryon density. As a result, its contribution predominantly occurs at early epochs and progressively vanishes with time, as realized also from the fact that below $z = 100$ the T_K curve is simply shifted to an higher adiabat. This fact causes the spin temperature to remain closer to (although always lower than) T_{CMB} , thus preserving the standard absorption feature. Quite remarkably the results assuming the realistic f_{abs} values do not differ appreciably from those obtained by imposing $f_{abs} = 1$ (Fig. 3, right panel). As pointed out before, the effects of annihilating DM are evident only at very high redshift, where the detailed calculation gives $f_{abs} \approx 1$ (see RFM06a), thus making the non-evolving f_{abs} approximation acceptable.

4.3 Global 21 cm background.

Having obtained the evolution of T_S for the different DM candidates, we are now ready to compute for each case the quantity that is most readily associated with observations, *i.e.* the differential brightness temperature, δT_b . These are shown in Fig. 4 for the realistic (left panel) and the constant f_{abs} (right panel) cases, respectively. In addition, to facilitate the comparison, we have also plotted in Fig. 5 the




Figure 4. *Left panel:* 21 cm differential brightness temperature as a function of redshift. The solid line shows δT_b without decaying/annihilating DM; while the long dashed, short dashed and dotted lines refer to δT_b with 25-keV decaying WDM, 10-MeV decaying LDM and 10-MeV annihilating LDM, respectively. Calculations were performed assuming f_{abs} as in RMF06a. *Right:* 21 cm differential brightness temperature as a function of redshift as in the left panel. Calculations were performed assuming $f_{abs} = 0.5$ for sterile neutrinos and $f_{abs} = 1$ in the other cases.

brightness temperature deviation $\delta T_b - \delta T_{b,0}$ of the various models from the standard one, restricted to the realistic f_{abs} evolution only.

As usual, we start our analysis from the realistic f_{abs} case. For sterile neutrinos (long-dashed curves) the characteristic absorption feature in the 21 cm background signal expected at $30 \leq z \leq 300$ is only slightly modified from the standard case (solid line): the maximum difference is found to be only ≈ -2 mK in the range $z \sim 10-40$. Yet, such a small signature of WDM decays could still be in principle observable: even modest sized single-dish radio telescopes can reach the required mK sensitivity in an all sky observation. The real challenge for the observation is represented by the ability to disentangle this cosmological signal from the various foregrounds (particularly the Galactic synchrotron emission) which could be several orders of magnitude brighter. The case of decaying 10-MeV LDM (short-dashed lines in Fig.4) is more interesting, as the difference with the standard case are larger. For this case we find that the deviation is predicted to be $\approx -(5-8)$ mK in the range $20 < z < 40$. Such an amplitude could be detected by LOFAR or SKA after a 1000 hour all sky integration and by most single dish radio telescopes, provide foreground contamination is taken care of. Finally, when LDM annihilations are considered, a considerably different δT_b evolution is obtained with respect to the standard case. Larger deviations are present in the entire range $40 < z < 200$, where the energy input of LDM annihilations forces δT_b to values larger than -20 mK, about two times smaller (in absolute value) than for the standard evolution. Such a large difference could in principle facilitate discriminating the annihilation scenario from the standard one. In practice, though, foregrounds and

ionospheric contamination become more severe at the low observing frequencies implied by these high redshifts.

Although of academic interest only, it is instructive to compare the constant f_{abs} cases (Fig. 4, right panel) with the previous results. As seen clearly from the Figure, considerably different conclusions would be drawn. First, sterile neutrino would drive δT_b to positive values, resulting in an emission signal below $z = 20$; a similar trend is also obtained for LDM decays, whose effects are seen in 21 cm emission with an amplitude of about 10 mK. The results for LDM annihilation instead do not show appreciable dependencies on the f_{abs} prescription, for the reasons discussed above.

5 DISCUSSION

From the results obtained in this paper, we conclude that it is in principle possible to observe the HI 21 cm signal from the Dark Ages produced by the energy input due to decays/annihilations of the most popular light/warm DM candidates. If so, radio observations might represent one of the most promising tools to study the nature of DM, as different particles are predicted to leave a specific signature on the signal. The sensitivity required to measure the 21 cm background signal can be achieved not only by the next generation of radio interferometers, but also by existing radio observatories.

However, the various foregrounds (*i.e.* Galactic free-free and synchrotron emission, unresolved extra-galactic radio sources, free-free emission from ionizing sources, synchrotron emission from cluster radio halos and relics) are much stronger than the cosmological signal, and will cer-

tainly prove extremely difficult to remove. Hence, a clear detection could be challenging to achieve.

It is beyond the scope of this paper to deal in detail with the problems of ionospheric scintillation and foreground contamination. A number of studies have discussed the foregrounds complications in some detail (e.g. Shaver et al. 1999; Oh & Mack 2003; Di Matteo, Ciardi & Miniati 2004) and we refer the reader to those papers for more information. For our aims it is sufficient to say that in general the sky temperature can be roughly described as:

$$T_{sky} \sim 180 \left(\frac{\nu}{180 \text{ MHz}} \right)^{-2.6} \text{ K} \quad (20)$$

(see e.g. Furlanetto, Oh & Briggs 2006) and is, at the frequencies of interest, several orders of magnitude higher than the cosmological 21 cm background signal.

The success of the 21 cm background observations to constrain DM will depend on the capability of effectively removing the foregrounds and of correcting for ionospheric variations (e.g. Gnedin & Shaver 2004; Zaldarriaga, Furlanetto & Hernquist 2004; Santos, Cooray & Knox 2005; Morales, Bowman & Hewitt 2005).

An alternative method to study the 21 cm backgrounds is by separating its spectral features from the smooth foreground spectrum (e.g. Furlanetto 2006, Shaver et al. 1999). Fig. 6 shows the gradient of δT_b for the DM candidates studied here. It is clear from the figure that the gradients are higher and easier to constrain in the frequency range $\nu = 1420/(1+z) \text{ MHz} \sim 10\text{--}40 \text{ MHz}$. The foreground gradients however increase with decreasing frequency making it difficult to predict the best frequencies to get to for a successful detection.

Discriminating among different DM cases would imply the ability to distinguish the difference in brightness temperature and/or in its gradient with respect to the standard scenario. We define these two quantities as $\Delta\delta T_b = \delta T_b - \delta T_b^0$ and $\Delta\delta T_b/\Delta\nu = d\delta T_b/d\nu - d\delta T_b^0/d\nu$, where the 0 stands for the standard (*i.e.* non decaying/annihilating) DM scenario.

Fig. 7 shows $\Delta\delta T_b$ versus $\Delta\delta T_b/\Delta\nu$ for the three DM models: the blue triangles correspond to 10-MeV annihilations, the red squares to 10-MeV decays and the black stars to the 25-keV sterile neutrino decays. The points are separated by 1 MHz steps in a range that goes from 5 to 130 MHz. As the sensitivity is somewhat uncertain for the different proposed 21 cm experiments, we assume as a guideline that a successful detection requires $\Delta\delta T_b > 3 \text{ mK}$ and $\Delta\delta T_b/\Delta\nu > 0.6 \text{ mK/MHz}$, respectively. The adopted value for $\Delta\delta T_b$ is a conservative one given that mK sensitivities are already achievable in all sky observations even by existing single dish radio telescopes. It has been chosen so to partially take into consideration the difficulties associated with foregrounds removal.

The shaded area covers the parameter space corresponding to likely non-detections. For the 25-keV sterile neutrinos case the points lie almost entirely within the shaded area, thus future observations would hardly be able to pin down this scenario.

In the case of decaying or annihilating 10-MeV LDM positive detections could be achieved on different frequency ranges. For the decays the ideal frequencies to study would be 40–80 MHz, where $\Delta\delta T_b \gtrsim 4 \text{ mK}$, and 80–90 MHz, where $\Delta\delta T_b \sim 2 \text{ mK}$ but $\Delta\delta T_b/\Delta\nu \gtrsim 0.6 \text{ mK/MHz}$. At




Figure 5. 21 cm differential brightness temperature difference in mK between the three DM cases considered and a non decaying–annihilating DM scenario. The dashed, solid and dotted lines correspond to 25-keV sterile neutrino decays, 10-MeV LDM decays and 10-MeV annihilations case respectively.

$\nu \geq 90 \text{ MHz}$, or $z \leq 15$ the first luminous sources start having a dominant impact on the IGM and the small modifications from decaying/annihilating DM would not be visible.

For the 10-MeV LDM annihilations case the best observing frequencies seem to be 10–30 MHz. At these frequencies $\Delta\delta T_b \gtrsim 6 \text{ mK}$ with a peak of over 20 mK at 10–20 MHz. Observations at such low frequencies are particularly difficult and probably future radio interferometers will not reach such large wavelengths. Probably 30 MHz is the minimum frequency which will be achieved by next generation instruments, and the frequency at which it will then be possible to constrain 10-MeV annihilating LDM.

We conclude then that it is likely that future observations of the 21 cm background radiation will allow us to effectively constrain among some of the DM candidates, which by decays or annihilations can influence deeply the properties of the IGM.

Nevertheless our study shows that accounting for the physical evolution of $f_{abs}(z)$ strongly reduces the amplitude of DM effects on the IGM with respect to those predicted by previous studies: for example we find that the effect of 25-keV decaying sterile neutrinos would be hardly observable.

REFERENCES

- Abazajian K., 2006, *PhRvD*, 73f, 3506
- Abazajian K., Fuller G. M., Tucker W. H., 2001, *ApJ*, 562, 593
- Abazajian K., Koushiappas S. M., 2006, *Phys.Rev.*, D74, 023527
- Allison, A. C., Dalgarno, A., 1969, *ApJ*, 158, 423
- Beacom J. F., Bell N. F., Bertone G., 2005, *PhRvL*, 94q, 1301
- Beacom J. F., Yüksel H., 2006, *PhRvL*, 97g, 1102




Figure 6. Gradient of the 21 cm differential brightness temperature. The solid line shows $d\delta T_b/d\nu$ without decaying/annihilating DM; while the long dashed, short dashed and dotted lines refer to $d\delta T_b/d\nu$ with 25-keV decaying WDM, 10-MeV decaying LDM and 10-MeV annihilating LDM, respectively. Calculations were performed assuming f_{abs} as in RMF06a.




Figure 7. Deviations of the differential brightness temperature (y -axis) and its gradient (x -axis) from those predicted in the standard case. The diagram illustrates the capability of constraining DM with future 21 cm observations. The black stars, red squares and blue triangles represent 25-keV decaying WDM, 10-MeV decaying LDM and 10-MeV annihilating LDM, respectively. The shaded area indicates the parameter space in which detection is excluded given the expected observational capabilities. Numbers along the curves refer to frequencies in MHz units.

Biermann P. L., Kusenkov A., 2006, PhRvL, 96L, 1301
 Bowman, J., Morales, M., Hewitt, J., 2005, astro-ph/0507357
 Boyarsky A., Neronov A., Ruchayskiy O., Shaposhnikov M., 2006a, MNRAS, 370, 213, astro-ph/0512509
 Boyarsky A., Neronov A., Ruchayskiy O., Shaposhnikov M., 2006b, astro-ph/0603368
 Chen, X. & Kamionkowski, M., 2004, PhRvD, 70, 043502
 Chen, X. & Miralda-Escudé, J. 2004, ApJ, 602, 1
 Ciardi, B. & Madau, P. 2003, ApJ, 596, 1
 Di Matteo, T., Ciardi, B., Miniati, F., 2004, MNRAS, 355, 1053
 Dolgov A. D., 2002, Phys.Rept., 370, 333
 Field, G. B. 1959, ApJ, 129, 551
 Furlanetto S. R., 2006, MNRAS, 371, 867
 Furlanetto S. R., Oh S. P., Briggs, 2006, PhR, 433, 181F
 Furlanetto S. R., Oh S. P., Pierpaoli E., 2006, PhRvD, 74j, 3502
 Gnedin, N. Y., Shaver, P. A., 2004, ApJ, 608, 611
 Hansen S. H., Haiman Z., 2004, ApJ, 600, 26
 Hirata, C.M. 2005, ApJ, in press (astro-ph/0507102)
 Kassim, N. E., Lazio, T. J. W., Ray, P. S., Crane, P. C., Hicks, B. C., Stewart, K. P., Cohen, A. S., & Lane, W. M. 2004, Planet. Space Sci., 52, 1343
 Knödseder J. et al., 2005, A&A, 441, 513
 Liszt, H., 2001, A&A, 371, 698L
 Madau, P., Meiksin, A., & Rees, M. J. 1997, ApJ, 475, 492
 Mapelli M., Ferrara A., 2005, MNRAS, 364, 2
 Mapelli M., Ferrara A. & Pierpaoli E., 2006, MNRAS, 369, 1719
 Morales, M. F., Bowman, J. D., Hewitt, J. N., 2005, astro-ph/0510027
 Oh, S. P., Mack, K. J., 2003, MNRAS, 346, 871
 Padmanabhan N., Finkbeiner D. P., 2005, PhRvD, 72b, 3508
 Pen, U. L., Wu, X. P., Peterson, J. 2004, astro-ph/0404083
 Pengelly, R. M., 1964, MNRAS, 127, 145
 Pierpaoli E., 2004, PhRvL, 92, 031301
 Ripamonti E., Mapelli M., Ferrara A., 2006a, MNRAS, accepted,

astro-ph/0606482 [RMF06a]
 Ripamonti E., Mapelli M., Ferrara A., 2006b, MNRAS, submitted, astro-ph/0606483 [RMF06b]
 Santos, M. G., Cooray, A., Knox, L., 2005, ApJ, 625, 575
 Seager, S., Sasselov, D. D., Scott, D., 1999, ApJ, 523, 1
 Shaver, P., Windhorst, R., Madau, P., de Bruyn, G., 1999, A&A, 345, 380
 Shchekinov Y.A., Vasiliev E.O., 2006, MNRAS, submitted, astro-ph/0604231
 Shull J. M., 1979, ApJ, 234, 761
 Shull J. M., van Steenberg M. E., 1985, ApJ, 298, 268
 Spergel, D. N. et al., 2006, ApJ, submitted
 Watson C. R., Beacom J. F., Yüksel H., Walker T. P., 2006, PhRvD, 74c, 3009
 Wouthuysen, S. A. 1952, AJ, 57, 31
 Wyithe, J.S., Loeb, A., Barnes, D., 2005, astro-ph/0506045
 Zaldarriaga, M., Furlanetto, S. R., Hernquist, L., 2004, ApJ, 608, 622
 Zhang L., Chen X., Lei Y.-A., Si Z., 2006, PhRvD, submitted, astro-ph/0603425